

テーマA: 「増殖シグナルを標的とした革新的がん治療法探索・開発」

- 1) Satoh R., Hagihara K., Matsuura K., Manse Y., Kita A., Kunoh T., Masuko T., Moriyama M., Moriyama H., Tanabe G., Muraoka O., Sugiura R.: Identification of ACA-28, a 1'-Acetoxychavicol Acetate analog compound, as a novel modulator of ERK MAPK signaling, which preferentially kills human melanoma cells. *Genes Cells*, in press (2017) 査読有り
- 2) Kobayashi A., Kanaba T., Satoh R., Ito Y., Sugiura R., Mishima M.: Chemical shift assignments of the first and second RRM of Nrd1, a fission yeast MAPK-target RNA binding protein. *Biomol NMR Assign.* (2017)
- 3) Chantarasrivong C., Ueki A., Ohyama R., Unga J., Nakamura S., Nakanishi I., Higuchi Y., Kawakami S., Ando H., Imamura A., Ishida H., Yamashita F., Kiso M., Hashida M.: Synthesis and functional characterization of novel sialyl LewisX mimic-decorated liposomes for E-selectin-mediated targeting to inflamed endothelial cells. *Mol. Pharmaceutics*, 24, in press. (2017) 査読有り
- 4) Nagai N., Ogata F., Deguchi S., Ueno A., Kawasaki N., Ito Y.: Combination ointment containing solid tranilast nanoparticles and dissolved sericin is efficacious for treating skin wound-healing deficits and redness in diabetic rats, *Biol. Pharm. Bull.* in press. (2017) 査読有り
- 5) Yamada H., Matsumura C., Yamada K., Teshima K., Hiroshima K., Kinoshita M., Suzuki S., Kakehi K.: Combination of SDS-PAGE and intact mass analysis for rapid determination of heterogeneities in monoclonal antibody therapeutics. *Electrophoresis*, in press 査読有り
- 6) * Yamagami M., Matsui Y., Hayakawa T., Yamamoto S., Kinoshita M., Suzuki S.: Plug-plug kinetic capillary electrophoresis for in-capillary exoglycosidase digestion as a profiling tool for the analysis of glycoprotein glycans. *J. Chromatogr. A*, in press. 査読有り
- 7) * Maegawa T., Nogata M., Honda T., Nakamura A., Miki Y.: 2-Arylquinoline synthesis from Cbz-protected 2-aminochalcone mediated by $\text{BF}_3 \cdot \text{Et}_2\text{O}$. *Heterocycles*, in press. (2017)
- 8) Ogata F., Nagai N., Kawasaki N.: Adsorption capability of cationic dyes (methylene blue and crystal violet) onto poly- γ -glutamic acid, *Chem. Pharm. Bull.* 65(3), 268-275 (2017) 査読有り
- 9) Ogata F., Toda M., Otani M., Kawasaki N.: Granulation of cobalt-containing nickel hydroxide with polyethylene terephthalate and its phosphate ion adsorption capability, *J. Water. Environ. Technol.* 15(1), 22-34 (2017) 査読有り
- 10) Ogata F., Ueta E., Toda M., Otani M., Kawasaki N.: Adsorption of phosphate ions from an aqueous solution by calcined nickel-cobalt binary hydroxide, *Water Sci. Technol.* 75(1), 94-105 (2017) 査読有り
- 11) * Ninomiya K., Chaipech S., Kunikata Y., Yagi R., Pongpiriyadacha Y., Muraoka O., Morikawa T.: Quantitative determination of stilbenoids and dihydroisocoumarins in *Shorea roxburghii* and evaluation of their hepatoprotective activity. *Int. J. Mol. Sci.*, 18, 451 (2017) 査読有り

- 12) Nagatomo Y., Hashimoto S., Kishimoto Y., Hayakawa T., Yamamoto S., Kinoshita M., Suzuki S.: Online cleanup liquid chromatography for the analysis of glycoprotein-derived oligosaccharides labeled with 7-amino-4-methylcoumarin. *Chromatography*, 38, 23-30 (2017) 査読有り
- 13) Okada M., Yamamoto A., Aizawa SI., Taga A., Terashima H., Kodama S. : HPLC Separation of Sulforaphane Enantiomers in Broccoli and Its Sprouts by Transformation into Diastereoisomers Using Derivatization with (S)-Leucine. *J Agric Food Chem.* 65, 244-250 (2017) 査読有り

テーマB: 「癌・増殖シグナル制御メカニズム解析」

- 1) Fukao A. and Fujiwara T. : The coupled- and uncoupled mechanisms by which trans-acting factors regulate mRNA stability and translation. *The Journal of Biochemistry.* (in press) 査読有り
- 2) Ogata F., Satoh R., Kita A., Sugiura R., Kawasaki N.: Evaluation of a novel method for measurement of intracellular calcium ion concentration in fission yeast. *J Toxicol Sci*, 42:159-166(2017)
- 3) Satoh R., Matsumura Y., Tanaka A., Takada M., Ito Y., Hagihara K., Inari M., Kita A., Fukao A., Fujiwara T., Hirai S., Tani T., Sugiura R.: Spatial regulation of the KH domain RNA-binding protein Rnc1 mediated by a Crm1-independent nuclear export system in *Schizosaccharomyces pombe*. *Mol Microbiol*, in press (2017)
- 4) * Kawase A., Hashimoto R., Shibata M., Shimada H., Iwaki M.: Involvement of reactive metabolites of diclofenac in diclofenac-induced cytotoxicity in sandwich-cultured rat hepatocytes. *Drug Metab Dispos*, 45, 316-324 in press. 査読有
- 5) * Tsubaki M., Takeda T., Kino T., Sakai K., Itoh T., Imano M., Nakayama T., Nishio K., Satoh T., Nishida S.: Contributions of MET activation to BCR-ABL1 tyrosine kinase inhibitor resistance in chronic myeloid leukemia cells. *Oncotarget.*, in press (2017). 査読有り
- 6) * Ota K., T Tong KI., Goto K., Tomida S., Komuro A., Wang Z., Nishio K., Okada H.: The H3K27 demethylase, Utx, regulates adipogenesis in a differentiation stage-dependent manner. *PLOS ONE* (in press) 査読有
- 7) Nishida S., Tsubaki M.: Exploration of Molecular Targets in the Development of New Therapeutics Aimed at Overcoming Multidrug Resistance. *Yakugaku Zasshi.*, 137, 145-149 (2017). 査読有り
- 8) Tsubaki M., Fujiwara D., Takeda T., Kino T., Tomonari Y., Itoh T., Imano M., Satou T., Sakaguchi K., Nishida S.: The sensitivity of head and neck carcinoma cells to statins is related to the expression of their Ras expression status, and statin-induced apoptosis is mediated via suppression of the Ras/ERK and Ras/mTOR pathways. *Clin Exp Pharmacol Physiol.*, 44, 222-234 (2017). 査読有り
- 9) Fujiwara D., Mashimo K., Kimura K., Noda A., Taki K., Yoshibayashi H., Takeda T., Tsubaki M., Nishida S., Sakaguchi K.: Influence of Next-Day Administration of Pegfilgrastim after FEC100 Chemotherapy in Japanese with Breast Cancer on Neutrophil Count. *Gan To Kagaku Ryoho.*, 44, 149-152 (2017) 査読有り

- 10) Tsubota M., Miyamoto T., Hiruma S., Saeki H., Miyazaki T., Sekiguchi F., Funakami Y., Kawabata A.: Repeated cold stress reduces cyclophosphamide-induced cystitis/bladder pain and macrophage activity in mice. *Pharmacology*, 99, 286-290 (2017) 査読有り
- 11) 坪田真帆、川畑篤史: トロンボモジュリンアルファの内臓痛治療への応用. *潰瘍*, in press (2017). 査読無し
- 12) * Ikuta H., Kawase A., Iwaki M.: Stereoselective pharmacokinetics and chiral inversion of ibuprofen in adjuvant-induced arthritic rats. *Drug Metab Dispos*, 45, 316-324 (2017)査読有
- 13) Fukami K., Asano E., Ueda M., Sekiguchi F., Yoshida S., Kawabata A.: High glucose induces N-linked glycosylation-mediated functional upregulation and overexpression of Cav3.2 T-type calcium channels in neuroendocrine-like differentiated human prostate cancer cells. *J. Pharmacol. Sci.*, 133, 57-60 (2017). 査読有り
- 14) Miyamoto T., Funakami Y., Kawashita E., Tomita S., Nomura A., Sugimoto N., Saeki H., Tsubota M., Ichida S., Kawabata A.: Enhanced hyperthermic responses to lipopolysaccharide in mice exposed to repeated cold stress. *Pharmacology*, 99, 172-178 (2017). 査読有り
- 15) Miyamoto T., Funakami Y., Kawashita E., Nomura A., Sugimoto N., Saeki H., Tsubota M., Ichida S., Kawabata A.: Repeated cold stress enhances the acute restraint stress-induced hyperthermia in mice. *Biol. Pharm. Bull.*, 40, 1-6 (2017). 査読有り
- 16) Terada Y., Tsubota M., Sugo H., Wakitani K., Sekiguchi F., Wada K., Takada M., Oita A., Kawabata A.: Tacrolimus triggers TRPV1-dependent relapse of pancreatitis-related pain in mice. *Pharmacology*, 99, 281-285 (2017). 査読有り
- 17) Terada Y., Wada K., Matsuda S., Kuwahara T., Kawabata A., Takada M., Watanabe T., Nakajima S., Sato T., Seguchi O., Yanase M., Fukushima N., Nakatani T.: Circadian pharmacokinetics and limited sampling strategy of everolimus in heat transplant patients. *Int. J. Clin. Pharmacol. Ther.*, 55, 1-8 (2017). 査読有り

テーマC: 「臨床評価・臨床腫瘍検体を用いた創薬シーズのPOC研究と個別化医療」

- 1) Mizukami T., Togashi Y., Naruki S., Banno E., Terashima M., de Velasco MA., Sakai K., Yoneshige A., Hayashi H., Fujita Y., Tomida S., Nakajima TE., Fujino T., Boku N., Ito A., Nakagawa K., Nishio K.: Significance of FGF9 gene in resistance to anti-EGFR therapies targeting colorectal cancer: A subset of colorectal cancer patients with FGF9 upregulation may be resistant to anti-EGFR therapies. *Mol Carcinog.* in press. 査読有り
- 2) Nakagawa K., Asamura H., Tsuta K., Nagai K., Yamada E., Ishi G., Mitsudomi T., Ito A., Higashiyama M., Tomita Y., Inoue M., Morii E., Matsuura N., Okumura M.: Novel method (OSNA) for the diagnosis of lymph node metastasis in patients with non-small cell lung cancer (NSCLC): Results of a multicenter prospective study. *Lung Cancer.* in press. 査読有り
- 3) Okada H., Tong K., Ota K., Komuro A., Ueda T., Ito A., Koch CA.: Attenuated DNA damage repair delays therapy-related myeloid neoplasms in a mouse model. *Cell Death Dis.* in press. 査読有り

- 4) Iino T., Hagiyaama M., Furuno T., Ito A., Hosokawa Y.: Time-course statistical evaluation of intercellular adhesion maturation by femtosecond laser impulse. *Biophys J.* in press. 査読有り
- 5) Yoneshige A., Hagiyaama M., Inoue T., Tanaka T., Ri A., Ito A.: Modest static pressure can cause enteric nerve degeneration through ectodomain shedding of cell adhesion molecule 1. *Mol Neurobiol.* in press. 査読有り
- 6) Tanizaki J., Hayashi H., Kimura M., Tanaka K., Takeda M., Shimizu S., Ito A., Nakagawa K.: Report of two cases of pseudoprogression in patients with non-small cell lung cancer treated with nivolumab – including histological analysis of one case after tumor regression. *Lung Cancer.* in press. 査読有り
- 7) Ohta M., Hosokawa Y., Hatano N., Sugano A., Ito A., Takaoka Y.: Efficacy of femtosecond lasers for application of acupuncture therapy. *Lasers in Med Sci.* in press. 査読有り
- 8) Kato T., Mizuguchi N., Ito A.: Candesartan suppresses proteinuria and decrease of nephrin expression but hydralazine does not in hypertensive nephropathy. *J Biomed.* in press. 査読有り
- 9) * Matsuoka H., Yoshiuchi K., Koyama A., Makimura C., Fujita Y., Tsurutani J., Sakai K., Sakamoto R., Nishio K., Nakagawa K.: Expectation of a Decrease in Pain Affects the Prognosis of Pain in Cancer Patients: a Prospective Cohort Study of Response to Morphine. *Int J Behav Med*, E-pub ahead of print, (2017) 査読有り
- 10) * Oiso N., Sakai K., Nishio K., Kawada A.: Phylloid hypomelanosis associated with a mosaic trisomy 13 in the 13q31.3-qter region: atypical phylloid distribution and typical hypomelanosis. *Pigment Cell Melanoma Res*, E-pub ahead of print, (2016) 査読有り
- 11) * Kobayashi Y., Azuma K., Nagai H., Kim YH., Togashi Y., Sesumi Y., Chiba M., Shimoji M., Sato K., Tomizawa K., Takemoto T., Nishio K., Mitsudomi T.: Characterization of EGFR T790M, L792F, and C797S mutations as mechanisms of acquired resistance to afatinib in lung cancer. *Mol Cancer Ther*, 16(2), 357-64 (2017) 査読有り
- 12) * Iwama E., Sakai K., Azuma K., Harada T., Harada D., Nosaki K., Hotta K., Ohyanagi F., Kurata T., Fukuhara T., Akamatsu H., Goto K., Shimose T., Kishimoto J., Nakanishi Y., Nishio K., Okamoto I.: Monitoring of somatic mutations in circulating cell-free DNA by digital PCR and next-generation sequencing during afatinib treatment in patients with lung adenocarcinoma positive for EGFR activating mutations. *Ann Oncol*, 28(1), 136-41 (2017) 査読有り